

Non-perturbative Renormalization Constants using Ward Identities

Tanmoy Bhattacharya,^a Shailesh Chandrasekharan,^b Rajan Gupta, Weonjong Lee, and Stephen Sharpe^{c,d}

^aMS B-285, Los Alamos National Lab, Los Alamos, New Mexico 87545, USA

^bDepartment of Physics, Duke University, Durham, North Carolina 27705, USA

^cPhysics Department, University of Washington, Seattle, Washington 98195, USA

^dCenter for Computational Physics, University of Tsukuba, Tsukuba, Ibaraki, 305-8577, Japan

We extend the application of axial Ward identities to calculate b_A , b_P and b_T , coefficients that give the mass dependence of the renormalization constants of the corresponding bilinear operators in the quenched theory. The extension relies on using operators with non-degenerate quark masses. It allows a complete determination of the $O(a)$ improvement coefficients for bilinears in the quenched approximation using Ward Identities alone. Only the scale dependent normalization constants Z_P^0 (or Z_S^0) and Z_T are undetermined. We present results of a pilot numerical study using hadronic correlators.

Wilson's discretization of the Dirac operator introduces lattice artifacts at $O(a)$, the effects of which are large for lattice spacings accessible in present simulations. It is thus expedient to devise improved lattice discretizations, and significant recent progress has been made in this direction. In particular the ALPHA collaboration has implemented Symanzik's improvement program [1], in which both the action and external sources are improved by the addition of higher dimensional operators. A key ingredient is the development of methods to determine the coefficients of these extra operators, the "improvement coefficients", non-perturbatively. In this note we present an extension of these methods which allows the determination of all improvement coefficients for bilinear operators.

We begin by reviewing the results of the ALPHA collaboration. They have shown that $O(a)$ artifacts can be removed from on-shell quantities by the addition of a single local operator of dimension five [2], resulting in the Sheikholeslami-Wohlert (or "clover") action [3]

$$S = S_G + S_W + S_{SW} \quad (1)$$

$$S_{SW} = a^5 c_{SW} \sum_x \bar{\psi}(x) \frac{i}{4} \sigma_{\mu\nu} F_{\mu\nu}(x) \psi(x). \quad (2)$$

Here S_G and S_W are respectively Wilson's gluon and quark actions, and S_{SW} is the Sheikholeslami-Wohlert term ($F_{\mu\nu}$ is the lattice gluon field strength tensor). Improvement of quark bilinears is slightly more involved [2]. We consider only flavor off-diagonal operators, the bare

lattice forms of which are

$$A_\mu^{(jk)} \equiv \bar{\psi}^j \gamma_\mu \gamma_5 \psi^k, \quad (3)$$

$$V_\mu^{(jk)} \equiv \bar{\psi}^j \gamma_\mu \psi^k, \quad (4)$$

$$P^{(jk)} \equiv \bar{\psi}^j \gamma_5 \psi^k, \quad (5)$$

$$S^{(jk)} \equiv \bar{\psi}^j \psi^k, \quad (6)$$

$$T_{\mu\nu}^{(jk)} \equiv \bar{\psi}^j i \sigma_{\mu\nu} \psi^k, \quad (7)$$

with $j \neq k$ being flavor indices, $\sigma_{\mu\nu} = [\gamma_\mu, \gamma_\nu]/2$, $\gamma_5 = \gamma_1 \gamma_2 \gamma_3 \gamma_4$, and the hermitian γ matrices satisfy $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}$. Removal of $O(a)$ errors from the on-shell matrix elements of these bilinears requires both the addition of extra operators (except for P and S),

$$(A_I)_\mu \equiv A_\mu + a c_A \partial_\mu P \quad (8)$$

$$(V_I)_\mu \equiv V_\mu + a c_V \partial_\nu T_{\mu\nu} \quad (9)$$

$$(T_I)_{\mu\nu} \equiv T_{\mu\nu} + a c_T (\partial_\mu V_\nu - \partial_\nu V_\mu). \quad (10)$$

and the introduction of mass dependence,

$$(X_R)^{(ij)} \equiv Z_X^0 (1 + b_X a m_{ij}) (X_I)^{(ij)}. \quad (11)$$

Here $X = A, V, P, S, T$, the Z_X^0 are renormalization constants in the chiral limit, and $m_{ij} \equiv (m_i + m_j)/2$ is the average bare quark mass.

Complete $O(a)$ improvement of matrix elements requires that the coefficients c_{SW} , c_X , and b_X , as well as the matching constants Z_X^0 , be determined non-

perturbatively.¹ Previous work has shown how the enforcement of axial and vector Ward identities (WI) allows one to determine Z_V^0 , Z_A^0 and Z_P^0/Z_S^0 [4], c_{SW} , c_A and b_V [5–7], c_V [8], c_T [13], and $b_P - b_A$ and b_S [14]. We discuss here an extension that yields b_A , b_P , and b_T .² This provides an alternative to the non-perturbative method proposed in Ref. [13] which uses the short-distance behavior of two point functions. The two remaining constants Z_P^0 (or Z_S^0) and Z_T^0 are scale and scheme dependent, and cannot be determined using WI.

It is important to note that the relations we derive do not extend directly to the unquenched theory, which requires additional improvement constants and a more complicated set of conditions as will be presented in [16].

We begin by recalling the ALPHA method for determining c_{SW} and c_A [6]. The improved axial current should satisfy

$$\partial_\mu (A_R)_{\mu}^{(ij)} = (m_i^R + m_j^R)(P_R)^{(ij)} + \mathcal{O}(a^2), \quad (12)$$

when inserted between on-shell states. Here m^R is the renormalized quark mass. It follows that the ratio

$$2\tilde{m}_{ij} \equiv \frac{\sum_{\vec{x}} \langle \partial_\mu [A_\mu + ac_A \partial_\mu P]^{(ij)}(\vec{x}, t) J^{(ji)}(0) \rangle}{\sum_{\vec{x}} \langle P^{(ij)}(\vec{x}, t) J^{(ji)}(0) \rangle} \quad (13)$$

$$= (m_i^R + m_j^R) \frac{Z_P^0(1 + b_P am_{ij})}{Z_A^0(1 + b_A am_{ij})}. \quad (14)$$

should be independent both of the choice of sources J and of the time t , as long as $t \neq 0$, up to corrections of $\mathcal{O}(a^2)$. This is to be achieved by simultaneously tuning c_{SW} and c_A . Our implementation of this condition differs from the Schrödinger functional method of Ref. [6] in that we use standard two-point correlation functions, with a variety of choices of sources. We also fix c_{SW} *a priori* and use the condition to determine only c_A .

It is convenient for the following discussion to introduce b coefficients defined using the masses $\tilde{m}_{ij}$, which we refer to as WI masses, rather than the bare quark mass, i.e.

$$(X_R)^{(ij)} \equiv Z_X^0(1 + \tilde{b}_X a \tilde{m}_{ij})(X_I)^{(ij)}. \quad (15)$$

The advantage of the WI mass is that it can be determined, using eq. (13), without the need for chiral extrapolation. By contrast, to determine the bare quark mass, $am = 1/(2\kappa) - 1/(2\kappa_c)$, one needs to know the critical hopping parameter, κ_c , the calculation of which requires chiral extrapolation. The difference is particularly significant when using standard two-point functions as opposed to the Schrödinger functional. To determine the standard b_X from the $\tilde{b}_X$ we need the relation between the bare

and WI masses, which is given in eq. (32) below. In the quenched theory it follows that

$$b_X = \frac{Z_P^0}{Z_A^0 Z_S^0} \tilde{b}_X + \mathcal{O}(a). \quad (16)$$

Note that we only need this relation at leading order in a , since the b_X appear only in $\mathcal{O}(a)$ corrections.

With c_{SW} fixed, Z_V^0 and $\tilde{b}_V$ can be obtained in the standard way using charge conservation. We use the forward matrix elements of $(V_I)_4$ between pseudoscalars,

$$\frac{1}{Z_V^0(1 + \tilde{b}_V a \tilde{m}_2)} = \frac{\sum_{\vec{x}, \vec{y}} \langle P^{(12)}(\vec{x}, \tau) (V_I)_4^{(22)}(\vec{y}, t) J^{(21)}(0) \rangle}{\langle \sum_{\vec{x}} P^{(12)}(\vec{x}, \tau) J^{(21)}(0) \rangle}. \quad (17)$$

with $\tau > t > 0$ and $J = P$ or A_4 . Here and below we use an abbreviation for the WI mass for two degenerate flavors: $\tilde{m}_i = \tilde{m}_{ii}$. Note that the c_V term in V_I does not contribute to the r.h.s., and so is not determined by this condition.

The remaining improvement constants are determined by enforcing the generic form of the integrated axial WI (AWI), in which the operator $\mathcal{O}^{23} = \bar{\psi}^{(2)} \Gamma \psi^{(3)}$ is transformed into $\delta \mathcal{O}^{13} = \bar{\psi}^{(1)} \gamma_5 \Gamma \psi^{(3)}$ by a chiral rotation in the 1, 2 flavor subspace:

$$\left\langle \delta \mathcal{S}^{(12)} \mathcal{O}_R^{(23)}(y) J^{(31)}(z) \right\rangle = \left\langle \delta \mathcal{O}_R^{(13)}(y) J^{(31)}(z) \right\rangle. \quad (18)$$

Here $\delta \mathcal{S}$ is the variation in the action

$$\delta \mathcal{S}^{(12)} = \int_{\mathcal{V}} d^4x \left[(m_1^R + m_2^R)(P_R)^{(12)} - \partial_\mu (A_R)_\mu^{(12)} \right], \quad (19)$$

and the chiral rotation is restricted to a 4-volume $\mathcal{V}$ containing y but not z . The Ward identities should hold up to corrections of $\mathcal{O}(a^2)$ if the operators and action are appropriately improved. There is, however, an obstacle to implementing these constraints, arising from the integral over x in $\delta \mathcal{S}$. This brings the pseudoscalar density $P_R(x)$ into contact with the operator $\mathcal{O}_R(y)$ on the l.h.s. of (18), implying that on-shell improvement of P and $\mathcal{O}$ is insufficient to improve the AWI. (The integral over $\partial_\mu A_\mu$ gives a surface term which does not involve contact with $\mathcal{O}$.) The problematic term is explicitly proportional to quark masses, and thus is absent in the chiral limit. For this reason the AWI has previously been used only in the chiral limit, from which one can determine Z_A^0 , Z_P^0/Z_S^0 , c_V and c_T , but not the $\tilde{b}_X$.

Our new observation is that by looking at the detailed dependence on the quark masses one can work away from the chiral limit and yet avoid contact terms. This allows

¹For brevity we refer to the Z_X^0 as improvement coefficients in the following.

²A preliminary account of this work was given in Ref. [15].

the determination of certain linear combinations of the $\tilde{b}_X$. The cost is the need to use non-degenerate quarks.

For all but the coefficient $\tilde{b}_T$, it is sufficient to work with two non-degenerate quarks: $m_1 = m_2 \neq m_3$. Since the contact term is explicitly proportional to $m_1^R + m_2^R$, which itself is proportional to $\tilde{m}_{12}$ at small quark masses, one can remove it by extrapolating to $\tilde{m}_1 = \tilde{m}_2 = 0$. The remaining $\tilde{m}_3$ dependence allows one to determine combinations of the $\tilde{b}_X$. In the following, we describe how this works for the different bilinears, and explain our particular implementation. The extrapolation to $\tilde{m}_1 = \tilde{m}_2 = 0$ will be implicit throughout this discussion.³

As a first application we show how to obtain $\tilde{b}_A - \tilde{b}_V$, as well as c_V , using the AWI with $\mathcal{O} = V_\mu$. The identity for the time component can be written

$$\begin{aligned} r_1 &\equiv \frac{\sum_{\vec{y}} \langle \delta \mathcal{S}_I^{(12)} (V_I)_4^{(23)}(\vec{y}, y_4) J^{(31)}(0) \rangle}{\sum_{\vec{y}} \langle (A_I)_4^{(13)}(\vec{y}, y_4) J^{(31)}(0) \rangle} \\ &= \frac{Z_A^0 (1 + \tilde{b}_A a \tilde{m}_3 / 2)}{Z_A^0 \cdot Z_V^0 (1 + \tilde{b}_V a \tilde{m}_3 / 2)}, \end{aligned} \quad (20)$$

where $\delta \mathcal{S}_I = - \int_V \partial_\mu (A_I)_\mu$. For the source we take $J = P$ or A_4 . We have chosen to put the current at $\vec{p} = 0$ so that the c_V term in V_I does not contribute. Since we know c_{SW} and c_A we can calculate r_1 , and determine $\tilde{b}_A - \tilde{b}_V$ from its dependence on $\tilde{m}_3$. The intercept provides a determination of Z_V^0 independent of that from eq. (17). The AWI for the spatial components can be written as

$$\begin{aligned} \tilde{r}_1 &= \frac{\sum_{\vec{y}} e^{i\vec{p} \cdot \vec{y}} \langle \delta \mathcal{S}_I^{(12)} [V_i + a c_V \partial_\mu T_{i\mu}]^{(23)}(\vec{y}, y_4) A_i^{(31)}(0) \rangle}{\sum_{\vec{y}} e^{i\vec{p} \cdot \vec{y}} \langle (A_I)_i^{(13)}(\vec{y}, y_4) A_i^{(31)}(0) \rangle}, \\ &= r_1. \end{aligned} \quad (21)$$

Enforcing this equality (for any small $\tilde{m}_3$) provides a determination of c_V .

Reversing the roles of vector and axial bilinears provides a second determination of $\tilde{b}_A - \tilde{b}_V$. For example, if c_V is known from above, then one can use the ratio

$$\begin{aligned} r_2 &\equiv \frac{\sum_{\vec{y}} e^{i\vec{p} \cdot \vec{y}} \langle \delta \mathcal{S}_I^{(12)} (A_I)_i^{(23)}(\vec{y}, y_4) V_i^{(31)}(0) \rangle}{\sum_{\vec{y}} e^{i\vec{p} \cdot \vec{y}} \langle (V_I)_i^{(13)}(\vec{y}, y_4) V_i^{(31)}(0) \rangle} \\ &= \frac{Z_V^0 (1 + \tilde{b}_V a \tilde{m}_3 / 2)}{Z_A^0 \cdot Z_A^0 (1 + \tilde{b}_A a \tilde{m}_3 / 2)}. \end{aligned} \quad (22)$$

Given Z_V^0 , this also yields Z_A^0 . The same information can be obtained from the combinations

$$\frac{1}{\sqrt{r_1 \cdot r_2}} = Z_A^0, \quad (23)$$

³ In practice we keep the term $\int (m_1 + m_2)(P_R)^{(12)}$ in eq. (18) prior to extrapolation, since this increases the range of $t_{\mathcal{O}}$ for which the ratio of correlation functions is constant.

$$\sqrt{\frac{r_1}{r_2}} = \frac{Z_A^0}{Z_V^0} (1 + (\tilde{b}_A - \tilde{b}_V) a \tilde{m}_3 / 2). \quad (24)$$

Applying the method to the AWI with $\mathcal{O} = P$ and S gives two determinations of $\tilde{b}_P - \tilde{b}_S$. For example, one can use the $\tilde{m}_3$ dependence of the ratio

$$\begin{aligned} r_3 &\equiv \frac{\sum_{\vec{y}} e^{i\vec{p} \cdot \vec{y}} \langle \delta \mathcal{S}_I^{(12)} S^{(23)}(\vec{y}, y_4) J^{(31)}(0) \rangle}{\sum_{\vec{y}} e^{i\vec{p} \cdot \vec{y}} \langle P^{(13)}(\vec{y}, y_4) J^{(31)}(0) \rangle} \\ &= \frac{Z_P^0 (1 + \tilde{b}_P a \tilde{m}_3 / 2)}{Z_A^0 \cdot Z_S^0 (1 + \tilde{b}_S a \tilde{m}_3 / 2)}, \end{aligned} \quad (25)$$

with $J = P$ or A_4 . Given Z_A^0 , this also yields Z_P^0 / Z_S^0 . Alternatively, one can use

$$\begin{aligned} r_4 &\equiv \frac{\sum_{\vec{y}} e^{i\vec{p} \cdot \vec{y}} \langle \delta \mathcal{S}_I^{(12)} P^{(23)}(\vec{y}, y_4) J^{(31)}(0) \rangle}{\sum_{\vec{y}} e^{i\vec{p} \cdot \vec{y}} \langle S^{(13)}(\vec{y}, y_4) J^{(31)}(0) \rangle} \\ &= \frac{Z_S^0 (1 + \tilde{b}_S a \tilde{m}_3 / 2)}{Z_A^0 \cdot Z_P^0 (1 + \tilde{b}_P a \tilde{m}_3 / 2)}, \end{aligned} \quad (26)$$

where for the source we choose either $J^{(31)} = S^{(31)}$ or $\sum_{\vec{z}} P^{(34)}(\vec{z}, z_4) P^{(41)}(0)$ for $0 < y_4 < z_4$.

The method described so far does not work for the tensor bilinear, because the chiral rotation transforms it back into (other components of) itself, and the dependence on $\tilde{b}_T$ cancels. One can, however, use the method to determine c_T [13]. For example, the AWI with $\mathcal{O} = T_{ij}$ at $\vec{p} = 0$ and $J = T_{k4}$, can be rearranged into

$$\begin{aligned} 1 + a c_T &\frac{\sum_{\vec{y}} \langle [-\partial_4 V_k]^{(13)}(\vec{y}, y_4) T_{k4}^{(31)}(0) \rangle}{\sum_{\vec{y}} \langle T_{k4}^{(13)}(\vec{y}, y_4) T_{k4}^{(31)}(0) \rangle} \\ &= Z_A^0 \frac{\sum_{\vec{y}} \langle \delta \mathcal{S}_I^{(12)} T_{ij}^{(23)}(\vec{y}, y_4) T_{k4}^{(31)}(0) \rangle}{\sum_{\vec{y}} \langle T_{k4}^{(13)}(\vec{y}, y_4) T_{k4}^{(31)}(0) \rangle}, \end{aligned} \quad (27)$$

Here we have moved the c_T dependence in $(T_I)_{k4}$ onto the l.h.s., and used the fact that $(T_I)_{ij}$ has no contribution from the c_T term at $\vec{p} = 0$. Given Z_A^0 , this equation determines c_T . A consistency check is that the result should be independent of $\tilde{m}_3$.

It turns out that one can determine b_T using the AWI (18), but to do so one must work with non-zero m_1 and m_2 . This means that the AWI is not completely improved: terms of $O(a)$ result from the contact of the pseudoscalar density in $\delta \mathcal{S}$ with the operator $\mathcal{O}$. The key point, however, is that these terms are proportional to $\tilde{m}_1 + \tilde{m}_2$, while the dependence on b_T is proportional to the difference $\tilde{m}_1 - \tilde{m}_2$. By separating these two dependences one can, in principle, determine b_T .

To explain this in detail we recall that off-shell $O(a)$ improvement requires the addition of an extra operator (multiplied by an extra improvement coefficient) for each

bilinear [13]. For the pseudoscalar and tensor the required additions are

$$P^{(12)} \longrightarrow P^{(12)} + ac'_P \bar{\psi}^{(1)}(-\overleftrightarrow{D} + m_1)\gamma_5 \psi^{(2)} + ac'_P \bar{\psi}^{(1)}\gamma_5(\overleftrightarrow{D} + m_2)\psi^{(2)}, \quad (28)$$

$$(T_I)^{(23)}_{\mu\nu} \longrightarrow (T_I)^{(23)}_{\mu\nu} + ac'_T \bar{\psi}^{(2)}(-\overleftrightarrow{D} + m_2)i\sigma_{\mu\nu}\psi^{(3)} + ac'_T \bar{\psi}^{(2)}i\sigma_{\mu\nu}(\overleftrightarrow{D} + m_3)\psi^{(3)}. \quad (29)$$

Inserting the off-shell improved operators into the AWI considered previously we find

$$r_5 \equiv \frac{\sum_{\vec{y}} e^{i\vec{p}\cdot\vec{y}} \langle \delta S_I^{(12)} (T_I)_{ij}^{(23)}(\vec{y}, y_4) T_{k4}^{(31)}(0) \rangle}{\sum_{\vec{y}} e^{i\vec{p}\cdot\vec{y}} \langle (T_I)_{k4}^{(13)}(\vec{y}, y_4) T_{k4}^{(31)}(0) \rangle} = \frac{1 + ab_T(\tilde{m}_1 - \tilde{m}_2)/2}{Z_A^0(1 - ab_A\tilde{m}_{12})} - 2a(c'_T + c'_P)\tilde{m}_{12}, \quad (30)$$

Note that both the $\tilde{b}_A$ term and the contact terms proportional to $c'_T + c'_P$ depend on $\tilde{m}_{12}$. Thus, assuming that c_T is known, $\tilde{b}_T$ can be obtained from the dependence of r_5 on $\tilde{m}_1 - \tilde{m}_2$. Note that the ratio r_5 depends on c_T at both zero and non-zero momentum.

A similar extension can be considered for the other AWI discussed above. It is straightforward to see that the $\tilde{m}_1 - \tilde{m}_2$ dependence of the ratios r_1 and r_2 allows a determination of $\tilde{b}_V + \tilde{b}_A$, while that of r_3 and r_4 gives a determination of $\tilde{b}_P + \tilde{b}_S$. Thus, in principle, one can determine all the $\tilde{b}_X$ using the generic AWI. One can also determine the five additional improvement constants c'_X using the dependence on $\tilde{m}_{12}$ as will be discussed in [16]. Further consistency checks are provided by the three-point vector WI with non-degenerate masses—although these by themselves do not allow one to disentangle the $\tilde{b}_X$ from the c'_X .

Petronzio and di Divitiis have shown that one can also use the two-point versions of the vector and axial WI to determine a subset of the quenched b_X , namely $b_A - b_P$, b_S and b_V [14]. The key point is again the use of non-degenerate quarks. In our numerical study we use some of their results, which we recall here.

The first result is obtained by comparing the WI mass $\tilde{m}$ to the bare quark mass m . These two masses are both related to the renormalized quark mass, the former through eq. (14), and the latter by

$$m_i^R = Z_m^0(1 + \tilde{b}_m a \tilde{m}_i) m_i \quad (31)$$

Combining these relations, and considering only degenerate masses $m_i = m_j$, one finds

$$\frac{m}{\tilde{m}} = \frac{Z_A^0}{Z_P^0 Z_m^0} [1 + (\tilde{b}_A - \tilde{b}_P - \tilde{b}_m) a \tilde{m}] \quad (32)$$

$$= \frac{Z_A^0 Z_S^0}{Z_P^0} [1 + (\tilde{b}_A - \tilde{b}_P + \tilde{b}_S/2) a \tilde{m}]. \quad (33)$$

To obtain the second line we have used the results $Z_m^0 Z_S^0 = 1$ and $b_S = -2b_m$, the latter valid only in quenched QCD [18]. Thus, from the slope and intercept of (33) one can determine $\tilde{b}_A - \tilde{b}_P + \tilde{b}_S/2$ and $Z_A^0 Z_S^0/Z_P^0$, respectively. To use this method we need to determine κ_c , which involves a chiral extrapolation.

The relationship between $\tilde{m}_{ij}$ and $(m_i + m_j)/2$ for non-degenerate quarks gives additional information. It turns out that this information can be gleaned without reference to the bare quark masses, and thus without the need for chiral extrapolation. In particular, by enforcing $(2m_1)^R + (2m_2)^R = 2(m_1 + m_2)^R$ to $O(a)$, one finds

$$\tilde{b}_P - \tilde{b}_A = -\frac{4\tilde{m}_{12} - 2[\tilde{m}_{11} + \tilde{m}_{22}]}{a[\tilde{m}_{11} - \tilde{m}_{22}]^2}. \quad (34)$$

This result, in terms of bare masses, was already noted in Ref. [14].

The final result from Ref. [14] uses the vector two-point WI. Requiring $\partial_\mu (V_R)_\mu^{(12)} = (m_1^R - m_2^R) S_R^{(12)}$ leads to

$$\Delta_{12} \equiv \frac{\sum_{\vec{x}} e^{i\vec{p}\cdot\vec{x}} \langle \partial_\mu V_{I\mu}^{(12)}(\vec{x}, t) J^{(21)}(0) \rangle}{\sum_{\vec{x}} e^{i\vec{p}\cdot\vec{x}} \langle S^{(12)}(\vec{x}, t) J^{(21)}(0) \rangle} \quad (35)$$

$$= \frac{Z_S[1 + \tilde{b}_S a \tilde{m}_{12}]}{Z_V[1 + \tilde{b}_V a \tilde{m}_{12}]} (m_1^R - m_2^R). \quad (36)$$

We implement this using two sources: $J^{(21)} = S^{(21)}$, and $J^{(21)} = \sum_{\vec{z}} P^{(23)}(\vec{z}, z_4) P^{(31)}(0)$ with $0 < t < z_4$. Enforcing the relation $(2m_1)^R - (2m_2)^R = 2(m_1 - m_2)^R$ to $O(a)$, with the l.h.s. determined from the AWI (14) and the rhs from the vector WI (35), we find

$$\frac{b_S - b_V}{2} + (b_P - b_A) = \frac{\Delta_{12} - R_Z[m_{11} - m_{22}]}{a R_Z[m_{11}^2 - m_{22}^2]}, \quad (37)$$

$$R_Z \equiv \frac{Z_S^0}{Z_P^0} \cdot \frac{Z_A^0}{Z_V^0}. \quad (38)$$

We can use eqs. (34,37) to determine b_P and b_S separately, since b_V and b_A are already known from eqs. (17,20).

We have performed a pilot test of our method on an ensemble of $16^3 \times 48$ quenched lattices at $\beta = 6/g^2 = 6.0$. We use the tree-level tadpole-improved value for the clover coefficient, $c_{SW} = 1.4755$, rather than the non-perturbative value $c_{SW} = 1.769$ [6]. This implies that our results for the b_X differ from the non-perturbative values by corrections of $O(g^2)$. We do not have data with three non-degenerate quarks, and so can test only the simpler version of our method which does not yield b_T . Nevertheless,

our results should suffice to assess the practicality of using WI with non-degenerate quarks to determine the b_X non-perturbatively.

Previous determinations of the improvement coefficients c_X have used Schrödinger functional boundary conditions in the time direction, with sources J placed on the boundaries. One advantage of this approach is that one can work directly in the chiral limit. In our study we use the same correlation functions as used in studying the spectrum and decay constants, i.e. we have periodic boundary conditions in the time direction. Thus a secondary output of our study is a comparison of these two approaches for determining the c_X . We stress, however, that our method for determining the b_X works for any choice of sources J , and in particular can be applied to the Schrödinger functional.

The correlation functions required in the integrated axial Ward identity (18) are obtained as follows. Quark propagators are calculated using a Wuppertal smeared source at $t = 0$ for five different values of κ corresponding to $aM_\pi = 0.57, 0.50, 0.43, 0.35, 0.24$. These propagators are used both to construct two-point functions and also as sources for propagators with the insertion of δS_I defined in eq. (19). Our insertion volume $\mathcal{V}$ is the region between $t = 4$ and 18. The second inversion uses the same κ , so that, as already noted, $m_1 = m_2$. To construct three-point functions, propagators with and without sources are contracted to form the operator $\mathcal{O}$. This allows us to insert any momentum into $\mathcal{O}$ and to place it anywhere in the interval $4 < t < 18$.

For the chiral extrapolations we ignore the correlations in the data between the different mass points. We find poor signals in some of the correlators containing quarks of the lightest mass, and so exclude the lightest mass from the extrapolations. The remaining four values of quark mass correspond to the range $m_s - 2.6m_s$, where m_s is the physical strange quark mass. Note that the relevant expansion parameter here is $m_q a$, and this is small for our range of quark masses.

Our results from the various WI are summarized in Table 1. Where there is a choice of sources, J , we have quoted the results with the best signal. In the following we discuss the various determinations pointing out salient features.

We determine c_A by requiring that the right hand side of eq. (13) is as close to a constant as possible over a range of times $t_{\min} - t_{\max}$. For long times only the pion contributes and the ratio is constant for all values of c_A —thus any choice of $t_{\max}$ in the pion-dominated region is equally good. To maximise our sensitivity to c_A , we choose $t_{\min}$ as small as possible while avoiding contact between δS_I and J . We find that our results are insensitive to small varia-

eq.	observable	two-point ∂	three-point ∂
13	c_A	-0.005(15)	-0.025(10)
17	Z_V^0	+0.745(1)	+0.746(1)
17	$\tilde{b}_V$	+1.57(2)	+1.57(2)
20	Z_V^0	+0.753(6)	+0.759(6)
20	$\tilde{b}_A - \tilde{b}_V$	-0.56(9)	-0.48(10)
21i	c_V	-0.66(27)	-0.65(25)
22i	$Z_V^0/(Z_A^0)^2$	+1.41(12)	+1.38(11)
22i	$\tilde{b}_A - \tilde{b}_V$	+1.2(1.2)	+1.8(1.0)
22ii	c_V	-0.30(26)	-0.31(24)
23	Z_A^0	+0.76(2)	+0.77(2)
24	Z_A^0/Z_V^0	+0.97(5)	+0.98(4)
24	$\tilde{b}_A - \tilde{b}_V$	+0.38(74)	+1.09(69)
25	$Z_A^0 Z_S^0/Z_P^0$	+0.95(2)	+0.98(1)
25i	$\tilde{b}_P - \tilde{b}_S$	-0.09(9)	-0.06(9)
27	c_T	+0.17(7)	+0.18(6)
33	$Z_A^0 Z_S^0/Z_P^0$	+0.97(1)	+0.97(1)
33	$\tilde{b}_A - \tilde{b}_P + \tilde{b}_S/2$	+0.48(1)	+0.50(1)
34	$\tilde{b}_P - \tilde{b}_A$	-0.20(9)	-0.13(8)
37	$\tilde{b}_S - \tilde{b}_V - 2\tilde{b}_P - 2\tilde{b}_A$	+0.04(48)	-0.27(38)

Table 1

Results for improvement coefficients from the listed WI. The two columns of results are for two choices of discretization of derivatives in eq. (13), as discussed in the text. The labels “i”–“v” are explained in the text.

tions in $t_{\min}$. There is a similar insensitivity to the choice of source J . This is in marked contrast to results from the unimproved action ($c_{SW} = 0$) [17] and gives us confidence that our tadpole-improved action has small enough $O(g^2 a)$ errors that we can carry out our tests.

To estimate the size of $O(a^2)$ errors (which become $O(a)$ errors in c_A), we use two choices of discrete derivatives in eq. (13): a scheme based on two-point derivatives,

$$\begin{aligned}\partial f(x+1/2) &= f(x+1) - f(x), \\ \partial^2 f(x+1/2) &= \frac{f(x+2) - f(x+1) - f(x) + f(x-1)}{2}, \\ f(x+1/2) &= \frac{f(x+1) + f(x)}{2},\end{aligned}$$

and one based on three-point derivatives,

$$\begin{aligned}\partial f(x) &= (f(x+1) - f(x-1))/2, \\ \partial^2 f(x) &= f(x+1) - 2f(x) + f(x-1).\end{aligned}$$

The $O(a^2)$ correction in the three-point ∂f is four times

larger than in the two-point case. The results for c_A from these two discretization schemes differ at $O(a)$.⁴ As shown in Table 1, this difference is substantial as, unfortunately, are the statistical errors. Because of this difference, we present results for the remaining improvement constants using both choices of c_A .

Both discretization schemes yield results for c_A significantly different from the value $c_A = -0.083(7)$ obtained previously at $c_{SW} = 1.769$ using the Schrödinger functional [6]. This difference is presumably a combination of $O(g^2)$ and $O(a)$ effects. We are unable to resolve this discrepancy in this work.

The signal in the vector WI (17) is very good, and provides our best estimates of Z_V and b_V . There is a tiny dependence on c_A due to the determination of the quark mass using eq. 13.

We now turn to results from our new method. We use a two-point discretization of the derivative in δS_I throughout.⁵ In Fig. 1 we show our results for the quantity r_1 which appears in the AWI (20). The data show a linear dependence on $\tilde{m}_3$ for our range of masses.

There is a numerical subtlety in the extraction of $\tilde{b}_A - \tilde{b}_V$. When using eq. 20, we have to make two choices concerning the form of $A_I^{(13)}$: whether to discretize it using two- or three-point derivatives; and whether to use the m dependent value of c_A obtained from eq. 13, or the chirally extrapolated value. Both choices only effect the result for $\tilde{b}_A - \tilde{b}_V$ at $O(a)$. In particular, it is straightforward to see that the two options for c_A lead to results for $\tilde{b}_A - \tilde{b}_V$ differing by $\sim aM_\pi^2/m$ (assuming pion domination of the correlators). Since M_π^2/m is a large scale, perhaps as large as 5 GeV, these differences, although technically of higher order, can be numerically significant. We find that they are only a 15% effect for the two-point discretization of A_I , but are much larger for the three-point discretization. For this reason we use the two-point discretization. As for the choice of c_A , we take the chirally extrapolated value, since this is the consistent choice at the order of improvement we are working. This does, however, have the disadvantage of giving poorer plateaus in the ratio r_1 .⁶

The results for Z_V^0 and $\tilde{b}_A - \tilde{b}_V$ are given in Table 1.

⁴Since our action is only perturbatively improved, one might worry that the two determinations of c_A could differ by $O(g^2)$ in addition to $O(a)$. This is not the case, however, because we use essentially the same matrix elements in both determinations, and so the difference between them is explicitly proportional to a .

⁵The results for improvement constants are expected to be fairly insensitive to this choice of discretization because, after integration over $\mathcal{V}$, different choices differ only by surface terms.

⁶We have a similar choice for the c_A appearing in $\delta S_I^{(12)}$, and here we use the mass dependent value which leads to better signals. This does not, however, directly affect our results for the $\tilde{b}_X$ since we extrapolate to $\tilde{m}_{12} = 0$.




Figure 1. The ratio r_1 as a function of $\tilde{m}_3/2$, after linear extrapolation to $\tilde{m}_1 = \tilde{m}_2 = 0$. The intercept and slope give $1/Z_V^0$ and $\tilde{b}_A - \tilde{b}_V$, respectively. The fit excludes the lightest point as the expected plateau in the fits to the different r_i are not uniformly good.

Z_V^0 is consistent, at the two σ level, with the result from the vector WI, but has much larger errors. $\tilde{b}_A - \tilde{b}_V$ is determined with a $\sim 20\%$ error.

The determination of c_V is more problematic and we have attempted a number of approaches. The two best are

- (i) We obtain c_V by first equating the right hand sides of Eqs. (20,21) for each $\tilde{m}_3$ and then extrapolating to $\tilde{m}_3 = 0$;
- (ii) For each $\tilde{m}_1 = \tilde{m}_2$ and $\tilde{m}_3$, we solve $r_1 r_2 = r_3 r_4$, where only r_2 depends on c_V . The chiral extrapolation in $\tilde{m}_1 = \tilde{m}_2$ removes the contribution of the contact terms. Lastly we extrapolate to $\tilde{m}_3 = 0$.

All methods require knowledge of c_A . The correction term proportional to c_V is $\approx 4\%$ in eq. (21) and $\approx 20\%$ in eq. (22), so one needs high statistics to get an accurate estimate. We prefer (i) as it is the most direct, and because r_4 does not have a good plateau. The large uncertainty in c_V extracted by these methods accounts for $\sim 50\%$ of the errors in Z_A^0 , and subsequently in Z_P^0/Z_S^0 and c_T .

With c_V (and Z_V^0) in hand, we use eq. (23) to determine our best estimate of Z_A^0 . The results from eqs. (22) and (24) are consistent but have larger errors. The estimates of $\tilde{b}_A - \tilde{b}_V$ from these two equations have very large errors.




Figure 2. The ratio r_3 as a function of $\tilde{m}_3/2$, after linear extrapolation to $\tilde{m}_1 = \tilde{m}_2 = 0$. The intercept and slope give $Z_P^0/(Z_S^0 Z_A^0)$ and $\tilde{b}_P - \tilde{b}_S$, respectively. The fit excludes the lightest point.

The final application of our new method is the determination of $\tilde{b}_P - \tilde{b}_S$ using eqs. (25) and (26). Note that neither of these requires knowledge of c_V . We find a good signal in the ratio r_3 but not in r_4 . The former is shown in Fig. 2 and the slope yields the value quoted in the table. The intercept gives an estimate of $Z_P^0/Z_S^0 Z_A^0$.

The last mixing coefficient c_T is determined from eq. (27). There is a good signal in all correlation functions and only c_A and Z_A^0 are needed beforehand. The error is dominated by the uncertainty in c_V which feeds in through Z_A^0 .

Further information, and consistency checks on the previous results, are provided by the WI using two-point functions. Equation (33) gives a result for $Z_P^0/Z_S^0 Z_A^0$, consistent with that from above, with similar errors. It also gives a very accurate result for the combination $\tilde{b}_A - \tilde{b}_P + \tilde{b}_S/2$, with little dependence on c_A . There is, however, an additional uncertainty due to the choice of κ_c . The two remaining equations, (34) and (37), which do not require κ_c , give rather poor determinations. An important technical point is that the $O(a)$ correction in eq. (35) is large when the source $J^{(21)}$ is chosen to be $S^{(21)}$ as $M_{\text{scalar}} a \approx 1$. In order to get results consistent with those from the choice $J^{(21)} = \sum_{\vec{z}} P^{(23)}(\vec{z}, z_4) P^{(31)}(0)$, we needed to use

	LANL	ALPHA	Pert. Th.
C_{SW}	1.4755	1.769	+1
Z_V^0	+0.745(1)(1)	0.7809(6)	+0.810
Z_A^0	+0.76(2)(1)	0.7906(94)	+0.829
Z_P^0/Z_S^0	+0.77(4)(1)	N.A.	+0.949
c_A	-0.005(15)(20)	-0.083(5)	-0.013
c_V	-0.66(27)(02)	-0.32(7)	-0.028
c_T	+0.17(7)(1)	N.A.	+0.020
$\tilde{b}_V$	+1.57(2)(1)	N.A.	+1.106
b_V	+1.62(3)(1)	1.54(2)	+1.273
$\tilde{b}_A - \tilde{b}_V$	-0.56(9)(8)	N.A.	-0.002
$\tilde{b}_P - \tilde{b}_S$	-0.09(9)(3)	N.A.	-0.066
$\tilde{b}_P - \tilde{b}_A$	-0.20(9)(7)	N.A.	+0.002
$\tilde{b}_A - \tilde{b}_P + \tilde{b}_S/2$	+0.48(1)(2)	N.A.	+0.584
$b_A - b_P + b_S/2$	+0.49(2)(2)	N.A.	+0.673
$\tilde{b}_A$	+1.01(09)(09)	N.A.	+1.104
$\tilde{b}_P$	+0.81(14)(16)	N.A.	+1.105
$\tilde{b}_S$	+0.90(17)(13)	N.A.	+1.172

Table 2

Our best estimates for normalization and improvement coefficients, compared to previous results where available, and to tadpole improved 1-loop perturbation theory (using $u_0 = 0.8778$). An estimate of the perturbative errors is $\alpha_s^2 \sim 0.02$. The first error is statistical, the second an estimate of the $O(a^2)$ error. See text for details.

a five-point discretization of $\partial_\mu V_\mu$ when using the source $J^{(21)} = S^{(21)}$.

We collect our best results in Table 2. These are obtained using the two-point derivative in δS_I . The difference between two- and three-point discretization is added as an additional error. Our results are compared to previous non-perturbative results, and to those of tadpole-improved 1-loop perturbation theory. The latter are obtained using 1-loop results available in the literature [9–12]. We note that for the c_X and the $\tilde{b}_X$ tadpole improvement is equivalent to using the boosted coupling g^2/u_0^4 in the 1-loop result. The same is not true for the Z_X and b_X . The conversion factor between the two definitions of the mass-dependent improvement coefficients, $\tilde{b}_X/b_X = Z_A^0 Z_S^0/Z_P^0$, turns out to be very close to unity (see Table 1). Thus, with the exception of b_V and $b_A - b_P + b_S/2$, which are determined quite accurately, we do not convert our results back to the standard definition b_X .

We draw the following conclusions. First, the method we have introduced appears to have practical utility. In the best channels, the statistical and systematic errors in the determination of the differences $\tilde{b}_A - \tilde{b}_V$ and $\tilde{b}_P - \tilde{b}_S$ are small compared to the values, $\tilde{b}_X \approx 1$, of the coefficients themselves. Second, c_V is determined rather poorly using our WI, and this accounts for a substantial fraction of the errors in Z_A^0 , Z_P^0/Z_S^0 , and c_T . For c_A and c_V , the Schrödinger functional method [5–8] gives results with much smaller errors. Third, we have found, in some cases, substantial disagreement between the perturbative predictions and our non-perturbative results. The most striking cases are b_V and $b_A - b_V$. These differences could be effects of $O(a^2)$, and we intend to investigate this issue by repeating the calculations at different values of a . Finally, there are also differences between our results and those of the ALPHA Collaboration. We anticipate that these differences are due in part to our use of an action that is not fully $O(a)$ improved. We have initiated simulations at $c_{SW} = 1.769$ to verify this.

We acknowledge the support of the Advanced Computing Laboratory at Los Alamos. This work was supported in part by the U.S. Department of Energy grants DE-LANL-ERWE161, DE-FG02-96ER40945, and DE-FG03-96ER40956. SS is very grateful to the Center for Computational Physics at the University of Tsukuba for the hospitality received there while completing this work.

REFERENCES

1. K. Symanzik, Nucl. Phys. **B226**, 187 (1983); p. 205 *ibid.*
2. K. Jansen *et al.*, Phys. Lett. **372B**, 275 (1996).
3. B. Sheikholeslami *et al.*, Nucl. Phys. **B259**, 572 (1985).
4. M. Bochicchio *et al.*, Nucl. Phys. **B262**, 331 (1985).
5. M. Lüscher *et al.*, Nucl. Phys. **B478**, 365 (1996).
6. M. Lüscher *et al.*, Nucl. Phys. **B491**, 323 (1997).
7. M. Lüscher *et al.*, Nucl. Phys. **B491**, 344 (1997).
8. M. Guagnelli and R. Sommer, Nucl. Phys. (Proc. Suppl.) **B63**, 886 (1998).
9. E. Gabrielli *et al.*, Nucl. Phys. **B362**, 475 (1991).
10. S. Sint *et al.*, Nucl. Phys. **B502**, 251 (1997).
11. S. Sint *et al.*, Nucl. Phys. (Proc. Suppl.) **B63**, 856 (1998).
12. Y. Taniguchi *et al.*, Phys. Rev. **D58**, 114503 (1998).
13. G. Martinelli *et al.*, Phys. Lett. **411B**, 141 (1997).
14. G.M. de Divitiis *et al.*, Phys. Lett. **419B**, 311 (1998).
15. T. Bhattacharya *et al.*, hep-lat/9810018.
16. T. Bhattacharya *et al.*, in preparation.
17. R. Gupta, hep-lat/9810016.
18. S. Sint and P. Weisz, Nucl. Phys. (Proc. Suppl.) **B63**,

856 (1998).